

KN-1

**BENZ- AND HETARENE-ANNULATED AZEPINES
FROM DONOR-ACCEPTOR CYCLOPROPANES**

I. V. Trushkov¹, A. E. Vartanova², A. Yu. Plodukhin³, O. A. Ivanova^{1,3}


¹ *N. D. Zelinsky Institute of Organic Chemistry, Russian Academy of Sciences, 47 Leninsky Pr., 119991, Moscow, Russia.*

² *Peoples' Friendship University of Russia, 6 Miklukho-Maklaya St., 117198, Moscow, Russia.*

³ *M. V. Lomonosov Moscow State University, 1-3 Leninskie Gory, 119991, Moscow, Russia.*

E-mail: itrushkov@mail.ru

Abstract. Donor-acceptor cyclopropanes are now considered as excellent building blocks for the synthesis of diverse carbo- and heterocycles including complex polycyclic structure. Herein, we report new methods for the synthesis of benz- and hetarene-annulated azepines based on two different approaches. The first one is based on the alkylation of 6-aminouracil and related substrates in *ortho*-position to the nucleophilic amino group followed by base-induced cyclization *via* attack of the amino group on the ester or keto group which was presented in the starting cyclopropane. The second one is based on the synthesis of *N*-aryl-1-methoxycarbonylcyclopropanecarboxamide bearing at the C(2) atom of the threemembered ring a donor substituent. Under the treatment with Lewis acid, these substrates undergo cyclization to 5-arylbenz[*b*]azepin-2-ones. The process diastereoselectivity was found to be efficiently controlled by the reaction time. Under short duration (1–2 h), *cis*-products were formed; the increase of the reaction time to 1 d led to the exclusive formation of *trans*-isomers.



Other transformations of donor-acceptor cyclopropanes to diverse heterocycles will be also discussed.

This work was supported by Russian Science Foundation (grant № 18-13-00449)